



Def: k -uniform $k \geq 2$ hypergraph $H = (V, E)$
 $V \equiv$ set of vertices $E \equiv$ set of edges
 $\forall e \in E$ is subset of V of size k .

Ex: $k=2$
graphs



$k=3$
Fano
plane



Remark: k -uniform hypergraph \approx pure (homogeneous) simplicial complex

Goal: hypergraphs are similar to graphs, yet many problems for hypergraph are exponentially harder. We illustrate this by surveying some open problems.

Ramsey th: Def: $r_k(s, t) \equiv \min N$ s.t. \forall red/blue coloring of ALL k -tuples on $[N]$ contains either red s -set or blue t -set. (Red-set \equiv all $\binom{[N]}{k}$ k -tuples are red)


If $s=t=n$ $r_k(s, t) = r_k(n)$. If $k=2$ we omit it.

Q: How large are $r_k(s, t)$. (finite by Ramsey th)

Th
E-S, E

$$2^{\lfloor n/2 \rfloor} \leq r(n) \leq 2^{2n}$$

- $r(s, t) \leq r(s-1, t) + r(s, t-1)$
- $N = 2^{\lfloor n/2 \rfloor}$. Colored edges R/B randomly
 $\Pr[\text{red/blue } n\text{-set}] \leq \binom{N}{n} \cdot 2 \cdot 2^{-\binom{N}{2}} \ll 1$



Th
AKS, K

$$r(3, t) = \Theta\left(\frac{t^2}{\log t}\right)$$

What about hypergraphs

$k=3$ Deleting one vertex gives
 $r_3(s, t) \leq r_2(r_3(s-1, t), r_3(s, t-1))$

Random coloring of triples shows that
 $r_3(n) \geq 2^{cn^2}$

Random coloring gives no bound on $r_3(4, n)$

Th:
E-R $r_3(s, t) \leq 2^{\binom{r(s-1, t-1)}{2}} \Rightarrow r_3(n) \leq 2^{2^{cn}}$

Problem 1: Show $r_3(n)$ is double exponential

Remark: $r_3(n, n, n)$ is! done by stepping up lemma.



Game: Players: Builder & Painter

Step $m+1$: new vertex v_{m+1} arrives. $\forall v_j, j \leq m$
 Builder decides if to add (v_j, v_{m+1}) . Once new edge is in
 Painter paints it R/B.

$r(s, t) \equiv$ min # of edges of Builder which force
 a red s -clique or blue t -clique

Th $r_3(s, t) \leq 2^{\tilde{r}(s, t)} \Rightarrow r_3(4, n) \leq 2^{cn \log n}$
 C-F-S

Lower Bound $r_3(4, n) \geq 2^{cn \log n}$ (CFS)

Take random tournament on $N = 2^{cn}$ vert. EASIER bound
 red edge  blue edge 

Problem: close gap $2^{cn \log n} \leq r_3(4, n) \leq 2^{cn^2 \log n}$

Discrepancy: Def: set is ALMOST MONOCHROM $\equiv (1-\epsilon)$ fraction
 of k -tuples have the same color.

FACTS: \forall R/B coloring of K_N has ALM. mon. set of size $O(\epsilon \log N)$

\forall R/B coloring of triples on $[N]$ has ALM. monoch. set of size $O(\epsilon \sqrt{\log N})$
 (C.F.S.)

NOTE: One would expect monochrom. set of size only $\log \log N$!


Problem: Prove \forall R/B coloring of k -tuples on $[N]$ has ALM. monochrom set of size $O(\epsilon \log^{\frac{1}{k-2}} N)$

Remark one expects $r_k(n) = 2^{2^{\dots^{2^n}}}$ tower.


Turan problem

Def: H -fixed k -uniform hypergraph $ex(n, H) \equiv$ max # of edges in n -vertex k -uniform hypergraph not containing H .

Th $ex(n, \Delta) = \frac{n^2}{4}$



Th $ex(n, K_{r+1}) = (1 - \frac{1}{r}) \frac{n^2}{2}$



Th
E-S $\Rightarrow \chi(H) = r+1$
(define chromatic)

Then $ex(n, H) = \left(1 - \frac{1}{r}\right) \frac{n^2}{2} + o(n^2)$


Min K_T s.t. vertices can be K -colored so that no edge is monochrom.

SOLUTION for ALL $r > 1$

Th
K-S-T
A-R-S

$ex(n, K_{s,t}) \geq cn^{2-1/s}$

, tight $t \geq (s-1)!$
in particular



sst

$ex(n, K_{2,2}) = cn^{3/2}$
 $ex(n, K_{3,3}) = cn^{5/3}$

construction
(n, d, λ) graphs
Define: Finite field
simple alg. geom


Hypergraphs

$K=3$ $K_4^3 \equiv 4$ vertices with ALL Triples

Q
Turán

$ex(n, K_4^3) \equiv ?$

Conj:



$n/3$

Def:

$\pi(H) = \lim_{n \rightarrow \infty} \frac{ex(n, H)}{\binom{n}{k}}$

Difficult

exponentially many non-isomorphic extremal constructions.

Conj: $\pi(K_4^3) = 5/9$

\$3000

Q:


what is $\pi(K_{k+1}^k)$?

Known:

$\frac{1}{k} \leq 1 - \pi(K_{k+1}^k) \leq \frac{\log k}{2k}$

min density in K -graph with no indep set of size $k+1$.

Q:



$K_{2,2,2}$

ALL triples intersecting
 v_1, v_2, v_3
 $|v_i|=2$

what is

$ex(n, K_{2,2,2}) \leq cn$

$3 - \frac{1}{2 \cdot 2}$

(maybe some construction of complexes with good eigenvalues might help)

Ruzsa-Szemerédi (Erdős-Brown-Sos)

Def: $f(n, p, q) = \max \#$ of triples in 3-graph s.t. no p vertices span $\geq q$ triples.

$f(n, 6, 3) = ?$

$(6, 3) \Rightarrow$ implies



no



no



$\Rightarrow f(n, 6, 3) \leq \frac{1}{3} \binom{n}{2}$


every triple covers 2 pairs
every pair covered once.

Th:
R-S

$$f(n, 6, 3) = o(n^2)$$

pf uses reduction to graphs and regularity lemma

R-S $\Rightarrow \forall A \subseteq [N]$
 $|A| = \delta N$ contains 3-term AP. Roth Th



$\forall x \in [N], a \in A \equiv$ 3 AP free

This 3-graph is $(6, 3)$ free
has $|A| \leq N$ triples

Q:

$$f(n, 7, 4) = o(n^2) ?$$

Endos-Frank-Rodl

Note
no
codegree 3



so $f(n, 7, 4) \leq \frac{2}{3} \binom{n}{2}$

Def: (i) cycle in graph

$$v_1 \dots v_k \text{ s.t. } (v_i, v_{i+1}) \equiv \text{edge}$$

(ii) cycle in hypergraph (Berge cycle)



vertices v_1, \dots, v_k and edges e_1, \dots, e_k s.t. $v_i \in e_i \cap e_{i+1}$ (addition mod k)

(iii) girth \equiv length of the shortest cycle.

Remind Def $\chi(G) = \dots!$

Remark: If Graph has large girth \Rightarrow locally looks like tree so is 2-colorable locally. What about globally.

Th \exists k -uniform hypergraph with large girth and large chrom. #.

Sketch: $k=3$ girth $\equiv t$ take edges rand. with prob.

$$p = n^{-2 + \frac{1}{t+1}}, \quad \# \text{ of cycles of length } \leq t \approx n^{2t} p^t \ll n \quad \text{easy to destroy deleting few vertices}$$

Union bound $\alpha(t) \approx n^{\frac{1}{2(t+1)}} \text{ polylog } n \Rightarrow \chi \geq n^{\frac{1}{2(t+1)} - o(1)} = k$

Cor: \exists 3-uniform hypergraph with girth t , $\chi = k$ and at most $k^{2(t+1)+o(1)}$ vertices

Q: find explicit construction

Remark: in case of graphs one can get such constructions from (n, d, λ) graphs one can get such in particular for const d girth is logarithm

Def: (n, d, λ) graph n -vertices d -regular all $|\lambda_i| \leq \lambda$ $i \geq 2$

Even simpler question

If $\chi(G) = s$ then $e(G) \geq \binom{s}{2}$ right. need edge between every 2-color classes.
right $G = K_s$

Open problem: $f(k, s) = \min \# \text{ of edges in } k\text{-uniform hypergraph with chromatic number } s.$

$k = \text{large}$
 $s = \text{fixed}$

$$\sqrt{\frac{k}{\log k}} 2^k \leq f(k, 3) \leq ck^2 2^k$$

minimal non 2-colorable hypergraph

Pf probabilistic: interesting to get explicit constructions.

$k = \text{fixed}$
 $s = \text{large}$

$$f(k, s) = \binom{(s-1)(k-1)+1}{k}$$

complete k -graph on that many vert. has chrom # s .

Alon: can improve this by roughly $\frac{\log k}{k}$ using turan #

lower bound $\approx \left[\frac{(s-1)(k-1)}{k} \right]^{k-1}$ gap $\exp(k)$

Def: (-2) girth = min $t \geq 4$ such that H has t vertices spanning $t-2$ edges.
combinatorial girth

Def: Steiner triple system \equiv 3-graph covering every pair of vertices once

STS exist if $2 | n-1$ $3 | \binom{n}{2}$.

Conj Erdos \exists STS with high (-2) -girth

easier Conj Is there a 3 uniform hypergraph with cn^2 edges (fixed) and arbitrarily high (-2) girth


Remark: (6,3) condition can be thought of (-3) girth so then we have $o(n^2)$ edges.

Def: mod 2-cycle in H is a collection of edges that cover every vertex even # of times

Remark: cycles in graph has this property.


claim: let H be 4-uniform (4 is just for convenience) hypergraph on n vertices which has $\geq \binom{n}{2}$ edges. Then it has mod 2-cycle of order $\log n$.

Pf: Define G - vertices $\binom{n}{2}$ pairs disjoint pairs adjacent if $(x_1, y_1, x_2, y_2) \equiv$ edge of H




Note every edge of H gives 3 edges of G .
so $e(G) \geq 3 \binom{n}{2} \Rightarrow G$ contains G' with min deg ≥ 3

G' has cycle of length $O(\log n)$



every level doubles compare to previous so this can continue only $O(\log n)$ steps

cycle in $G \equiv$ mod 2-cycle in H



Conj: suppose H has $\geq \binom{n}{2}$ edges. Show still that it has mod 2 cycle of length polylog n.

Hamiltonicity


Def: Hamilton cycle is a cycle passing through every vertex of graph exactly once.

Q: When G is Hamiltonian (NP hard, suff. cond)

Expansion and HC

Posa: if $\forall x \subseteq G, |x| \leq k \implies |N(x)| \geq 2|x| \implies G$ has cycle of length $\geq 2k$

ROTATION



Th $K-S$ Let G be (n, d, λ) graph with $\lambda \leq \frac{d}{\log n}$ then G is Hamiltonian

Conj
 $\lambda \leq \frac{d}{c}$
 $c \equiv$ large constant is enough

Remark: since $\lambda \approx \sqrt{d}$ in best case works for $d \approx \log^2 n$ very sparse.

Def: H - 3 uniform hypergraph, n vert.

Tight HC ordering $v_1 \dots v_n$ s.t. $(v_i, v_{i+1}, v_{i+2}) \in \text{edge}$
 loose HC edges overlap in one vertex

Q: Find which spectral condition on H implies tight/loose HC