

Random Simplicial Complexes

Roy Meshulam

Technion – Israel Institute of Technology

Borel Seminar "High-Dimensional Expanders"

Les Diablerets, March 2015

Lecture I: Topology of the Binomial Model

Random Graphs

- ▶ The Erdős-Rényi model
- ▶ Topology of random graphs

The k -dimensional Erdős-Rényi model

- ▶ Homological connectivity
- ▶ Fundamental group
- ▶ The random k -tree process
- ▶ Collapsibility and top homology
- ▶ Embedding dimension

The Erdős-Rényi Model

$G(n, p)$ = random graphs on the $[n] = \{1, \dots, n\}$
with independent edge probabilities p .

Ubiquity of Random Graphs

- ▶ Models for a multitude of natural phenomena, e.g. phase transition problems of statistical physics.
- ▶ A useful tool for showing the existence (indeed the prevalence) of graphs with favorable properties that are hard (or perhaps impossible) to produce explicitly.

A Classical Example: Ramsey Graphs

Erdős ('48): A typical $G \in G(n, \frac{1}{2})$ will contain neither a **clique** nor an **independent set** on $k = 2 \log_2 n$ vertices.

Remark: Current best explicit constructions achieve only $k = 2^{(\log n)^\epsilon}$ (for any fixed $\epsilon > 0$).

Topology of a Random Graph

Theorem [Erdős-Rényi '58]:

For any function $\omega(n)$ that tends to infinity

$$\lim_{n \rightarrow \infty} \Pr [G \in G(n, p) : G \text{ connected}] =$$

$$\begin{cases} 0 & p = \frac{\log n - \omega(n)}{n} \\ 1 & p = \frac{\log n + \omega(n)}{n} \end{cases} .$$

$$\lim_{n \rightarrow \infty} \Pr [G \in G(n, \frac{c}{n}) : G \text{ acyclic}] =$$

$$\begin{cases} 0 & c > 1 \\ \sqrt{1-c} \cdot e^{\frac{2c+c^2}{4}} & c < 1 \end{cases} .$$

A Model of Random Complexes

Y a simplicial complex , $Y^{(i)} = i$ -dim skeleton of Y .

$Y^{(i)}$ = oriented i -dim simplices of Y .

$f_i(Y) = |Y^{(i)}|$.

Δ_{n-1} = the $(n - 1)$ -dimensional simplex on $V = [n]$.

$Y_k(n, p)$ = probability space of all complexes

$$\Delta_{n-1}^{(k-1)} \subset Y \subset \Delta_{n-1}^{(k)}$$

with probability distribution

$$\Pr(Y) = p^{f_k(Y)}(1 - p)^{\binom{n}{k+1} - f_k(Y)} .$$

Homological Connectivity of Random Complexes

Fix $k \geq 1$ and a finite abelian group R .

Theorem [Linial-M '03 , M-Wallach '06]:

For any function $\omega(n)$ that tends to infinity

$$\lim_{n \rightarrow \infty} \Pr [Y \in Y_k(n, p) : \tilde{H}_{k-1}(Y; R) = 0] = \begin{cases} 0 & p = \frac{k \log n - \omega(n)}{n} \\ 1 & p = \frac{k \log n + \omega(n)}{n} \end{cases} .$$

Theorem [Hoffman-Kahle-Paquette '13]:


If $p > \frac{80k \log n}{n}$ then a.a.s. $\tilde{H}_{k-1}(Y; \mathbb{Z}) = 0$.

Homological Connectivity - the Lower Bound

$\sigma \in \Delta_{n-1}(k-1)$ is **isolated** in $Y \in Y_k(n, p)$ if it is not contained in any $\tau \in Y(k)$.

$\sigma \in \Delta_{n-1}(k-1)$ isolated in Y

$$\Rightarrow 0 \neq 1_\sigma \in \tilde{H}^{k-1}(Y)$$



If $p = \frac{k \log n - \omega(n)}{n}$ then

$$E[\text{number of isolated } \sigma' \text{'s}] = \binom{n}{k} (1-p)^{n-k} \rightarrow \infty.$$

A second moment argument then implies that

$$\Pr[\tilde{H}_{k-1}(Y; R) \neq 0] \rightarrow 1.$$

Fundamental Group of Random 2-Complexes

Theorem [Babson-Hoffman-Kahle '10]:

For any $\epsilon > 0$

$$\Pr [Y \in Y_2(n, p) : \pi_1(Y) = \{1\}] \rightarrow \begin{cases} 0 & p = n^{-\frac{1}{2}-\epsilon} \\ 1 & p = n^{-\frac{1}{2}+\epsilon} \end{cases} .$$

Theorem [M '11]:

If $\{G_n\}$ is a sequence of finite groups with $|G_n| \leq n^c$ then


$$\Pr [Y \in Y_2(n, \frac{(3c+6) \log n}{n}) : \text{Hom}(\pi_1(Y), G_n) = \{1\}] \rightarrow 1.$$

Simple Connectivity - the Upper Bound

For $Y \in Y_2(n, p)$ and $u \in [n]$ let $Y_u = \text{St}_Y u$.

$$Y_u \cap Y_v =$$

$$\text{St}_Y uv \cup G$$



It follows that if $p \geq \sqrt{\frac{10 \log n}{n}}$ then $Y_u \cap Y_v$ is connected for all u, v . Therefore, by the nerve lemma, $\pi_1(Y) = \{1\}$.

Hyperbolic Groups

Finitely presented groups

X - finite set, $F(X)$ - free group on X .

$R \subset F(X)$ - finite set of relations.

$G = \langle X | R \rangle = F(X) / (\text{normal subgroup generated by } R)$.

Area

Let $w \in F(X)$ such that $\bar{w} = 1 \in G$.

$\text{area}(w) = \text{minimal } n \text{ such that } w = (u_1^{-1} r_1^{\epsilon_1} u_1) \cdots (u_n^{-1} r_n^{\epsilon_n} u_n)$

for some $u_i \in F(X), r_i \in R, \epsilon_i \in \{\pm 1\}$.

Hyperbolicity

G is hyperbolic if there exist a constant A such that

$$\text{area}(w) \leq A \cdot \text{length}(w)$$

for all $w \in F(X)$ with $\bar{w} = 1$.

Examples

Hyperbolic groups

- ▶ Finite groups.
- ▶ Free groups.
- ▶ Fundamental groups of surfaces of genus ≥ 2 :

$$G = \langle a_1, b_1, \dots, a_g, b_g \mid [a_1, b_1] \cdots [a_g, b_g] \rangle.$$


\mathbb{Z}^2 is not hyperbolic

$$\mathbb{Z}^2 = \langle a, b \mid aba^{-1}b^{-1} \rangle.$$

$$w_n = a^n b^n a^{-n} b^{-n}.$$

$$\text{length}(w_n) = 4n.$$

$$\text{area}(w_n) = \Omega(n^2).$$



Isoperimetry and Hyperbolicity

X - a 2-dimensional simplicial complex.

γ - a simplicial null-homotopic loop in X of length $|\gamma|$.

$A_X(\gamma)$ - the minimal number of simplicies in a filling of γ .

The Isoperimetric Constant

$$I(X) = \inf \left\{ \frac{|\gamma|}{A_X(\gamma)} : \gamma \sim 1 \right\}.$$

Theorem:

$$I(X) > 0 \iff \pi_1(X) \text{ Hyperbolic} .$$

Local to Global Principle

Theorem [Gromov]:

Let $c > 0$ and let X be a 2-dimensional complex that satisfies:

$$I(S) \geq c \text{ for all pure } S \subset X \text{ such that } f_2(S) \leq 10^6 c^{-2}.$$

Then:

$$I(X) \geq 10^{-2} c.$$

The Structure of Sparse Complexes

A Density Invariant

$$\tilde{\mu}(X) = \min \left\{ \frac{f_0(Z)}{f_2(Z)} : Z \subset X \right\}.$$

Theorem [Babson-Hoffman-Kahle]:

- ▶ $\tilde{\mu}(X) > \frac{1}{2} \Rightarrow X$ is homotopic to a wedge of circles, 2-spheres and projective planes.
- ▶ For any $\epsilon > 0$ there exists a $c_\epsilon > 0$ such that

$$\tilde{\mu}(X) > \frac{1}{2} + \epsilon \Rightarrow I(X) > c_\epsilon .$$

Simple Connectivity - the Lower Bound

Theorem [Babson-Hoffman-Kahle]:

$$p = o(n^{-\frac{1}{2}-\epsilon}) \Rightarrow \Pr[I(Y) > 0] = 1 - o(1).$$

Sketch of Proof: Let S be a pure complex such that $f_2(S) \leq 10^6 c_\epsilon^{-2}$. If $\tilde{\mu}(S) \leq \frac{1}{2} + \epsilon$ then there exists a subcomplex $Z \subset S$ such that $f_0(Z) \leq (\frac{1}{2} + \epsilon)f_2(Z)$ and hence

$$\Pr[Y \supset S] \leq n^{f_0(Z)} p^{f_2(Z)} = n^{(\frac{1}{2}+\epsilon)f_2(Z)} \cdot o(n^{-\frac{1}{2}-\epsilon})^{f_2(Z)} = o(1).$$

Therefore Y a.a.s. satisfies the following condition:

$$S \subset Y \text{ and } f_2(S) \leq 10^6 c_\epsilon^{-2} \Rightarrow \tilde{\mu}(S) > \frac{1}{2} + \epsilon \Rightarrow I(S) > c_\epsilon.$$


The local to global principle now implies that $I(Y) > 10^{-2} c_\epsilon$.

Structure of $\pi_1(Y)$

Theorem [Costa-Farber '13]:

- ▶ If $p = n^{-\frac{3}{5} + \epsilon}$ then a.a.s. $\pi_1(Y)$ has 2-torsion.
- ▶ If $p = n^{-\frac{3}{5} - \epsilon}$ then a.a.s. $\pi_1(Y)$ is torsion free.

The 6-Point Projective Plane \mathbb{P}^2



If $p = n^{-\frac{f_0}{2} + \epsilon} = n^{-\frac{3}{5} + \epsilon}$ then
a.a.s. $Y \supset Z \cong \mathbb{P}^2$ such that
 $\mathbb{Z}_2 = \pi_1(Z) \hookrightarrow \pi_1(Y)$.

Aspherical Subcomplexes in $Y_2(n, p)$

A 2-dimensional complex X is aspherical if $\pi_2(X) = 0$.

Whitehead Conjecture

Let X be 2-dimensional complex. If X is aspherical and $Z \subset X$, then Z is aspherical.

Theorem [Costa-Farber '13]:

If $p = n^{-\frac{1}{2}-\epsilon}$ then a.a.s. Y satisfies the Whitehead conjecture, i.e. If $Y' \subset Y$ is aspherical then so are all $Y'' \subset Y'$.

An Easy Upper Bound on the Threshold for $H_k(Y) = 0$

Let $z^{k-1}(Y) = \dim Z^{k-1}(Y)$, $\beta_k(Y) = \dim H_k(Y)$.

By Euler-Poincaré:

$$\beta_k(Y) = f_k(Y) - \binom{n}{k} + z^{k-1}(Y).$$

In particular, for $p = \frac{c}{n}$:

$$E[\beta_k] \geq E[f_k] - \binom{n-1}{k} \gtrsim \left(\frac{c}{k+1} - 1\right) \frac{n^k}{k!}.$$

Hence


$$c > k + 1 \Rightarrow \beta_k(Y) > 0 \text{ a.a.s.}$$

This can be (slightly) improved ...

k -Collapsibility

Suppose $\sigma \in X(k-1)$ is contained in a **unique** $\tau \in X(k)$.

An **Elementary k -Collapse**: $X \rightarrow X - \{\sigma, \tau\}$.




X is **k -Collapsible** if there is a sequence of k -collapses:

$$X = X_1 \rightarrow X_2 \rightarrow \cdots \rightarrow X_m$$

such that $\dim X_m \leq k-1$.

X k -collapsible $\not\Rightarrow$ X homotopic to a $(k-1)$ -complex

The Dunce Hat



A contractible but non-collapsible complex

Boundaries of $(k + 1)$ -simplices

A Trivial Obstruction

If $Y \supset \partial\Delta_{k+1}$ then $H_k(Y) \neq 0$ and hence Y is not k -collapsible.

The Distribution of Boundaries

$f(Y) =$ number of $\partial\Delta_{k+1}$ in $Y \in Y_k(n, \frac{c}{n})$.

$$E[f] = \binom{n}{k+2} p^{k+2} \sim \frac{c^{k+2}}{(k+2)!}.$$

$$f(Y) \sim \text{Poisson}(\lambda) \quad \text{with} \quad \lambda = \frac{c^{k+2}}{(k+2)!}.$$

The Threshold for k -Collapsibility

$x_k =$ minimal positive root of $\exp(-\frac{1-x}{kx}) = x$.

$\gamma_k = (kx_k(1-x_k)^{k-1})^{-1}$.

Remark: $\gamma_2 \doteq 2.455$, $\gamma_3 \doteq 3.089$, $\gamma_k = \Theta(\log k)$.

Theorem [Aronshtam-Linial-Łuczak-M]:

For any $c < \gamma_k$

$$\lim_{n \rightarrow \infty} \Pr [Y \in Y_k(n, \frac{c}{n}) : Y \text{ is } k\text{-collapsible}] =$$

$$\lim_{n \rightarrow \infty} \Pr [Y \not\supseteq \partial\Delta_{k+1}] = \exp(-\frac{c^{k+2}}{(k+2)!}).$$

Theorem [Aronshtam-Linial]:

For any $c > \gamma_k$

$$\lim_{n \rightarrow \infty} \Pr [Y \in Y_k(n, \frac{c}{n}) : Y \text{ is } k\text{-collapsible}] = 0.$$

The Threshold for $H_k(X) = 0$

y_k = minimal positive root of $\exp(-\frac{(k+1)(1-y)}{1+ky}) = y$.

$$\gamma_k^* = (k+1)(1+ky_k)^{-1}(1-y_k)^{-k}.$$

Remark: $\gamma_2^* \doteq 2.753$, $\gamma_3^* \doteq 3.907$, $\gamma_k^* = k+1 - \Theta(k^2 \exp(-k))$.

Theorem [Linial-Peled]:

For any $c < \gamma_k^*$

$$\lim_{n \rightarrow \infty} \Pr [Y \in Y_k(n, \frac{c}{n}) : H_k(Y; \mathbb{Q}) = 0] =$$

$$\lim_{n \rightarrow \infty} \Pr [Y \not\supseteq \partial \Delta_{k+1}] = \exp(-\frac{c^{k+2}}{(k+2)!}).$$

Theorem [Aronshtam-Linial]:


For any $c > \gamma_k^*$

$$\lim_{n \rightarrow \infty} \Pr [Y \in Y_k(n, \frac{c}{n}) : H_k(Y; \mathbb{Z}) = 0] = 0.$$

Rooted k -Trees

A k -Tree is a k -dimensional simplicial complex T on $V = \{v_1, \dots, v_\ell\}$ such that for all $k + 1 \leq i \leq \ell$ $\text{link}(X[v_1, \dots, v_{i-1}], v_i)$ is a $(k - 1)$ -simplex.
The **Root** of T is $\eta = \{v_1, \dots, v_k\}$.

2-Tree



The Random k -Tree Process [ALLM]

Fix k and a parameter $\gamma > 0$. Inductively construct a chain of **random** k -trees rooted at $\eta = \{12 \dots k\}$:

$$\{\eta\} = T_k(0, \gamma) \subset T_k(1, \gamma) \subset T_k(2, \gamma) \subset \dots$$

Suppose $T_k(r, \gamma)$ is given. **Independently** for each $(k-1)$ -face


$$\tau \in T_k(r, \gamma) - T_k(r-1, \gamma)$$

pick $J_\tau \sim \text{Poisson}(\gamma)$ new vertices $z_{\tau 1}, \dots, z_{\tau J_\tau}$ and let

$$T_k(r+1, \gamma) = T_k(r, \gamma) \cup \bigcup_{\tau} \{z_{\tau 1}\tau, \dots, z_{\tau J_\tau}\tau\}.$$

Fact: $T_k(r, \gamma)$ approximates the r -neighborhood of τ in $Y_k(n, \frac{\gamma}{n})$.

A 2-Tree Process



Pruning a Rooted k -Tree

Let T be a k -tree rooted at η .

$\{\tau_1, \dots, \tau_t\} = \text{Free } (k-1)\text{-faces of } T \text{ different from } \eta$.

$\sigma_i = \text{the unique } k\text{-simplex containing } \tau_i$.

A Pruning Step

$$T \rightarrow T - \{\tau_1, \sigma_1, \dots, \tau_t, \sigma_t\} .$$

Main Point

Let $\gamma_k = \text{maximal such that for } \gamma < \gamma_k$

$$\lim_{r \rightarrow \infty} \Pr[T_k(r, \gamma) \searrow \eta \text{ in } < r \text{ pruning steps}] = 1.$$

Then γ_k/n is the threshold for k -collapsibility.

Pruning of a Random k -Tree I

Probability of an Early Collapse

$$\rho_k(r, \gamma) = \Pr[T_k(r, \gamma) \searrow \eta \text{ in } < r \text{ pruning steps}].$$

A Recursion for $\rho_k(r, \gamma)$

$$\rho_k(1, \gamma) = \Pr[T_k(1, \gamma) = \eta] = \exp(-\gamma) .$$

$T_k(r, \gamma)$ collapses to η in $< r$ steps **iff** each k -simplex $\sigma \supset \eta$ contains a face $\eta' \neq \eta$ such that the random k -tree rooted in η' collapses to η' in $< r - 1$ steps. Therefore:

$$\rho_k(r, \gamma) = \sum_{j=0}^{\infty} \Pr [J = j] (1 - (1 - \rho_k(r - 1, \gamma))^k)^j = \exp(-\gamma(1 - \rho_k(r - 1, \gamma))^k) .$$

Pruning of a Random k -Tree II

Let


$$\rho_k(\gamma) = \lim_{r \rightarrow \infty} \rho_k(r, \gamma).$$

$\rho_k(\gamma)$ is the smallest positive root of the equation

$$u_k(\gamma, x) = \exp(-\gamma(1-x)^k) - x = 0.$$

Graphs of $u_2(\gamma, x)$ for

$\gamma = 0, 1, 2, 2.455$



Pruning of a Random k -Tree III

If $\gamma \geq 0$ is small then $\rho_k(\gamma) = 1$. Let

$$\gamma_k = \inf\{\gamma > 0 : \rho_k(\gamma) < 1\} \quad \text{and} \quad x_k = \rho_k(\gamma_k).$$

The pair (γ_k, x_k) satisfies

$$u_k(\gamma, x) = \frac{\partial u_k}{\partial x}(\gamma, x) = 0.$$

It follows that x_k is the unique solution of

$$\exp\left(-\frac{1-x}{kx}\right) = x$$

and

$$\gamma_k = (kx_k(1-x_k)^{k-1})^{-1}.$$


Embedding Dimension

Any k -dimensional complex can be embedded in \mathbb{R}^{2k+1} .


Examples of k -dimensional complexes not embeddable in \mathbb{R}^{2k} :

The k -skeleton of the $(2k + 2)$ -simplex.

The $(k + 1)$ -fold join of 3-point spaces.



K_5



$K_{3,3}$

Theorem [Wagner]:

There exist constants $c_k < C_k$ such that:

- ▶ $Y \in Y_k(n, \frac{c_k}{n})$ a.a.s. embeds in \mathbb{R}^{2k} .
- ▶ $Y \in Y_k(n, \frac{C_k}{n})$ a.a.s. does not embed in \mathbb{R}^{2k} .

Lecture II: Cohomological Expansion

Expansion in Graphs and Complexes

- ▶ Cheeger constant of a graph
- ▶ Cohomological expansion of complexes
- ▶ Application to random complexes
- ▶ Expansion and topological overlap

Randomized construction of expanding complexes


- ▶ Expander graphs and complexes
- ▶ A bounded degree model
- ▶ Spectral gaps and 2-Expansion
- ▶ Random LS-Complexes are 2-Expanders

The Graphical Cheeger Constant

Edge Cuts

For a graph $G = (V, E)$ and $S \subset V$, $\bar{S} = V - S$ let

$$e(S, \bar{S}) = |\{e \in E : |e \cap S| = 1\}|.$$



Cheeger Constant

$$h(G) = \min_{0 < |S| \leq \frac{|V|}{2}} \frac{e(S, \bar{S})}{|S|}.$$

Spectral Gap

Laplacian Matrix

$G = (V, E)$ a graph, $|V| = n$.

The **Laplacian** of G is the $V \times V$ matrix L_G :

$$L_G(u, v) = \begin{cases} \deg(u) & u = v \\ -1 & uv \in E \\ 0 & \text{otherwise.} \end{cases}$$

Eigenvalues of L_G

$$0 = \mu_1(G) \leq \mu_2(G) \leq \dots \leq \mu_n(G).$$

$\mu_2(G) =$ **Spectral Gap** of G .

Expansion and Spectral Gap

Theorem [Alon-Milman, Tanner]:

For all $\emptyset \neq S \subsetneq V$

$$e(S, \bar{S}) \geq \frac{|S||\bar{S}|}{n} \mu_2.$$

In particular

$$h(G) \geq \frac{\mu_2}{2}.$$

Theorem [Alon, Dodziuk]:

If G is d -regular then

$$h(G) \leq \sqrt{2d\mu_2}.$$

Expanders can thus be defined using the spectral gap.

Simplicial Cohomology

X a simplicial complex on V , R a fixed abelian group.

i -face of $\sigma = [v_0, \dots, v_k]$ is $\sigma_i = [v_0, \dots, \widehat{v}_i, \dots, v_k]$.

$C^k(X) = k$ -cochains = skew-symmetric maps $\phi : X(k) \rightarrow R$.

Coboundary Operator $d_k : C^k(X) \rightarrow C^{k+1}(X)$ given by

$$d_k \phi(\sigma) = \sum_{i=0}^{k+1} (-1)^i \phi(\sigma_i) .$$

$d_{-1} : C^{-1}(X) = R \rightarrow C^0(X)$ given by

$d_{-1} a(v) = a$ for $a \in R$, $v \in V$.

$Z^k(X) = k$ -cocycles = $\ker(d_k)$.

$B^k(X) = k$ -coboundaries = $\text{Im}(d_{k-1})$.

k -th reduced cohomology group of X :

$$\tilde{H}^k(X) = \tilde{H}^k(X; R) = Z^k(X)/B^k(X) .$$


Cut of a Cochain

Cut determined by a k -cochain $\phi \in C^k(X; R)$:

$$\text{supp}(d_k\phi) = \{\tau \in X(k+1) : d_k\phi(\tau) \neq 0\}.$$

Cut Size of ϕ : $\|d_k\phi\| = |\text{supp}(d_k\phi)|$.

Example:




$$\|d_1\phi\| = |\{\sigma_1, \sigma_2\}| = 2$$

Hamming Weight of a Cochain

The **Weight** of a k -cochain $\phi \in C^k(X; R)$:

$$\|\phi\| = \min \{ |\text{supp}(\phi + d_{k-1}\psi)| : \psi \in C^{k-1}(X; R) \}.$$

Example: $\|\phi\| = 3$ but $\|[\phi]\| = 1$



Expansion of a Complex

Expansion of a Cochain

The expansion of $\phi \in C^k(X; R) - B^k(X; R)$ is

$$\frac{\|d_k \phi\|}{\|[\phi]\|}.$$

k -expansion Constant

$$h_k(X; R) = \min \left\{ \frac{\|d_k \phi\|}{\|[\phi]\|} : \phi \in C^k(X; R) - B^k(X; R) \right\}.$$

Remarks:

- ▶ $h_k(X; R) > 0 \Leftrightarrow \tilde{H}^k(X; R) = 0$.
- ▶ In the sequel: $h_k(X) = h_k(X; \mathbb{F}_2)$.

Expansion of a Simplex I

Δ_{n-1} = the $(n - 1)$ -dimensional simplex on $V = [n]$.

Claim [M-Wallach, Gromov]:

$$h_{k-1}(\Delta_{n-1}) = \frac{n}{k+1}.$$


Example:

$$[n] = \cup_{i=0}^k V_i, \quad |V_i| = \frac{n}{k+1}$$

$$\phi = \mathbf{1}_{V_0 \times \dots \times V_{k-1}}$$

$$\|\phi\| = \left(\frac{n}{k+1}\right)^k$$

$$\|d_{k-1}\phi\| = \left(\frac{n}{k+1}\right)^{k+1}$$



Expansion of a Simplex II

Let $\phi \in C^{k-1}(\Delta_{n-1})$. For $u \in V$ define $\phi_u \in C^{k-2}(\Delta_{n-1})$ by

$$\phi_u(\tau) = \begin{cases} \phi(u\tau) & u \notin \tau \\ 0 & u \in \tau \end{cases} .$$

Let $\sigma \in \Delta_{n-1}(k-1)$ and $u \in V$. Then

$$d_{k-1}\phi(u\sigma) = \phi(\sigma) - \sum_{w \in \sigma} \phi(u(\sigma - w)) = \phi(\sigma) - d_{k-2}\phi_u(\sigma)$$

Therefore

$$\begin{aligned} (k+1)\|d_{k-1}\phi\| &= |\{(\tau, u) \in \Delta_{n-1}(k) \times V : u \in \tau \in \text{supp}(d_{k-1}\phi)\}| \\ &= |\{(\sigma, u) \in \Delta_{n-1}(k-1) \times V : \sigma \in \text{supp}(\phi - d_{k-2}\phi_u)\}| \\ &= \sum_{u \in V} |\text{supp}(\phi - d_{k-2}\phi_u)| \geq n\|\phi\|. \end{aligned}$$

Homological Connectivity of Random Complexes

Fix $k \geq 1$ and a finite abelian group R .

Theorem [Linial-M '03 , M-Wallach '06]:

For any function $\omega(n)$ that tends to infinity

$$\lim_{n \rightarrow \infty} \Pr [Y \in Y_k(n, p) : \tilde{H}_{k-1}(Y; R) = 0] = \begin{cases} 0 & p = \frac{k \log n - \omega(n)}{n} \\ 1 & p = \frac{k \log n + \omega(n)}{n} \end{cases} .$$

The Relevance of Expansion:

If $0 \neq [\phi] \in \tilde{H}^{k-1}(\Delta_{n-1}^{(k-1)})$ then

$$\begin{aligned} \Pr [[\phi] \in \tilde{H}^{k-1}(Y; \mathbb{F}_2)] &= (1 - p)^{\|d_k \phi\|} \\ &\leq (1 - p)^{\frac{n \|\phi\|}{k+1}} . \end{aligned}$$

Expansion and Homological Connectivity

A weak threshold:

If $p = \frac{(k^2+k+1) \log n}{n}$ then a.a.s. $H^{k-1}(Y; \mathbb{F}_2) = 0$.

Proof:

$$\Pr [\tilde{H}^{k-1}(Y; \mathbb{F}_2) \neq 0] \leq \sum_{0 \neq [\phi] \in \tilde{H}^{k-1}(\Delta_{n-1}^{(k-1)})} (1-p)^{\|d_{k-1}\phi\|} \leq$$

$$\sum_{m \geq 1} \binom{\binom{n}{k}}{m} (1-p)^{\frac{nm}{k+1}} \leq$$

$$\sum_{m \geq 1} (n^k n^{-\frac{k^2+k+1}{k+1}})^m =$$

$$\sum_{m \geq 1} (n^{-\frac{1}{k+1}})^m \rightarrow 0.$$

Weighted Expansion

X - n -dimensional pure simplicial complex.

A probability distribution on $X(k)$:

$$w(\sigma) = \frac{|\{\eta \in X(n) : \sigma \subset \eta\}|}{\binom{n+1}{k+1} f_n(X)}.$$

For $\phi \in C^k(X)$ let

$$\|\phi\|_w = \sum_{\{\sigma \in X(k) : \phi(\sigma) \neq 0\}} w(\sigma)$$

$$\|[\phi]\|_w = \min\{\|\phi + d_{k-1}\psi\| : \psi \in C^{k-1}(X)\}.$$

Weighted k -th Expansion:

$$\underline{h}_k(X) = \min \left\{ \frac{\|d_k \phi\|_w}{\|[\phi]\|_w} : \phi \in C^k(X) - B^k(X) \right\}.$$

The Affine Overlap Property

Number of Intersecting Simplices


For $A = \{a_1, \dots, a_n\} \subset \mathbb{R}^k$ and $p \in \mathbb{R}^k$ let

$$\gamma_A(p) = |\{\sigma \subset [n] : |\sigma| = k + 1, p \in \text{conv}\{a_i\}_{i \in \sigma}\}|.$$

Theorem [Bárány]:

There exists $p \in \mathbb{R}^k$ such that

$$f_A(p) \geq \frac{1}{(k+1)^k} \binom{n}{k+1} - O(n^k).$$



The Topological Overlap Property

Number of Intersecting Images


For a continuous map $f : \Delta_{n-1} \rightarrow \mathbb{R}^k$ and $p \in \mathbb{R}^k$ let

$$\gamma_f(p) = |\{\sigma \in \Delta_{n-1}(k) : p \in f(\sigma)\}|.$$

Theorem [Gromov]:

There exists $p \in \mathbb{R}^k$ such that

$$\gamma_f(p) \geq \frac{2k}{(k+1)!(k+1)} \binom{n}{k+1} - O(n^k).$$



Topological Overlap and Expansion

Number of Intersecting Images

For a continuous map $f : X \rightarrow \mathbb{R}^k$ and $p \in \mathbb{R}^k$ let

$$\gamma_f(p) = |\{\sigma \in X(k) : p \in f(\sigma)\}|.$$

Expansion Condition on X

Suppose that for all $0 \leq i \leq k - 1$

$$\underline{h}_i(X) \geq \epsilon.$$

Theorem [Gromov]

There exists a $\delta = \delta(k, \epsilon)$ such that for any continuous map $f : X \rightarrow \mathbb{R}^k$ there exists a $p \in \mathbb{R}^k$ such that

$$\gamma_f(p) \geq \delta f_k(X).$$

Symmetric Matroids

Matroid:

An n -dimensional simplicial complex $M \subset 2^V$ such that $M[S]$ is pure for all $S \subset V$.

Homology of matroids:

$\tilde{H}_i(M) = 0$ for all $0 \leq i \leq \dim M - 1$.

Symmetric matroid:

$G = \text{Aut}(M)$ is transitive on the maximal faces.

Examples of Symmetric Matroids

The Partition Matroid $X_{n,m}$

Let V_1, \dots, V_{n+1} be $n + 1$ disjoint sets, $|V_i| = m$.

$$X_{n,m} = \left\{ \sigma \subset \bigcup_{i=1}^{n+1} V_i : \forall i \quad |\sigma \cap V_i| \leq 1 \right\}.$$

Independence Matroid of Affine Space

$$\text{IN}(\mathbb{F}_q^n) = \{ \sigma \subset \mathbb{F}_q^n : \sigma \text{ is linearly independent} \}.$$

Hermitian Unital with 65 Points

Independence matroid of the curve

$$H = \{ [x, y, z] \in PG(2, 16) : x^5 + y^5 + z^5 = 0 \}.$$

Expansion of Symmetric Matroids

Proposition [Lubotzky-M-Mozes]:

M symmetric matroid $\Rightarrow \underline{h}_k(M) \geq 8^{-\dim M} \quad \forall k \leq \dim M - 1.$

Example: The Partition Matroid $X_{n,m}$

For $0 \leq k \leq n - 1$

$$\underline{h}_k(X_{n,m}) \geq \frac{\binom{n+1}{k+1}}{\sum_{j=0}^{k+1} \left(\frac{2(m-1)}{m}\right)^j \binom{n-j}{n-k-1}}.$$

Corollary [Dotterrer-Kahle]:

$$\underline{h}_k(\text{Octahedral } n\text{-sphere}) = \underline{h}_k(X_{n,2}) \geq 1$$

and

$$\underline{h}_{n-1}(X_{n,m}) \geq \frac{n+1}{2^{n+1} - 1}.$$

Expansion of Spherical Buildings

The Building $A_{n+1}(\mathbb{F}_q)$

Vertices: All nontrivial linear subspaces $0 \neq V \subsetneq \mathbb{F}_q^{n+2}$.

Simplices: $V_0 \subset \cdots \subset V_k$.

Theorem [Solomon, Tits]:

$$\dim \tilde{H}_i(A_{n+1}(\mathbb{F}_q)) = \begin{cases} q^{\binom{n+2}{2}} & i = n \\ 0 & \text{else.} \end{cases}$$

Proposition [Gromov, LMM]:

$$\underline{h}_{n-1}(A_{n+1}(\mathbb{F}_q)) \geq \frac{1}{(n+2)!}.$$

Expander Graphs

(d, ϵ) -Expanders

A family of graphs $\{G_n = (V_n, E_n)\}_n$ with $|V_n| \rightarrow \infty$ with two seemingly contradicting properties:

- ▶ **High Connectivity:** $h(G_n) \geq \epsilon$.
- ▶ **Sparsity:** $\max_v \deg_{G_n}(v) \leq d$.

Pinsker:

Random $3 \leq d$ -regular graphs are (d, ϵ) -expanders.

Margulis:

Explicit construction of expanders.

Lubotzky-Phillips-Sarnak, Margulis:

Ramanujan Graphs - an "optimal" family of expanders.

The Ubiquity of Expanding Graphs

Uses of Expanders

- ▶ Construction of efficient communication networks.
- ▶ Randomization reduction in probabilistic algorithms.
- ▶ Construction of good error correcting (LDPC) codes.
- ▶ Tools in computational complexity lower bounds.

Interactions with Other Areas

- ▶ Expansion and Kazhdan's property T.
- ▶ Expanders as spaces of maximal Euclidean distortion.
- ▶ Dimension expanders and representation theory.
- ▶ Expanders on finite simple groups.

Expander Complexes

Degree of a Simplex

For $\sigma \in X(k-1)$ let $\deg(\sigma) = |\{\tau \in X(k) : \sigma \subset \tau\}|$.

$$D_{k-1}(X) = \max_{\sigma \in X(k-1)} \deg(\sigma).$$

(k, d, ϵ) -Expanders

A family of Complexes $\{X_n\}_n$ with $f_0(X_n) \rightarrow \infty$ such that

$$D_{k-1}(X_n) \leq d \quad \text{and} \quad h_{k-1}(X_n) \geq \epsilon.$$

Random Complexes as Expanders

$Y \in Y_k(n, p = \frac{k^2 \log n}{n})$ is a.a.s. a $(k, \log n, 1)$ -expander.

Problem

Do there exist (k, d, ϵ) -expanders with $k \geq 2$ and **fixed** d, ϵ ?

Latin Squares

Definitions

$\mathbb{S}_n =$ Symmetric group on $[n]$.

$(\pi_1, \dots, \pi_k) \in \mathbb{S}_n^k$ is **legal** if $\pi_i(\ell) \neq \pi_j(\ell)$ for all ℓ and $i \neq j$.

A **Latin Square** is a legal n -tuple $L = (\pi_1, \dots, \pi_n) \in \mathbb{S}_n^n$.

$\mathcal{L}_n =$ Latin squares of order n with uniform measure.

The Usual Picture

$L = (\pi_1, \dots, \pi_n) \leftrightarrow T_L \in M_{n \times n}([n])$

$T_L(i, \pi_k(i)) = k$ for $1 \leq i, k \leq n$.

Example for $n = 4$

$$\pi = (1234)$$

$$L = (Id, \pi, \pi^2, \pi^3)$$


$$T_L =$$

1	2	3	4
4	1	2	3
3	4	1	2
2	3	4	1

The Complete 3-Partite Complex

$$V_1 = \{a_1, \dots, a_n\}, \quad V_2 = \{b_1, \dots, b_n\}, \quad V_3 = \{c_1, \dots, c_n\}$$

$$T_n = V_1 * V_2 * V_3 = \{\sigma \subset V : |\sigma \cap V_i| \leq 1 \text{ for } 1 \leq i \leq 3\}$$



$$T_n \simeq S^2 \vee \dots \vee S^2 \quad (n-1)^3 \text{ times}$$

Latin Square Complexes


$L = (\pi_1, \dots, \pi_n) \in \mathcal{L}_n$ defines a complex $Y(L) \subset T_n$ by

$$Y(L)(2) = \{ [a_i, b_j, c_{\pi_i(j)}] : 1 \leq i, j \leq n \}.$$

Example: $n = 2$

$$L = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & 1 \\ \hline \end{array}$$

$$Y(L) =$$



$$Y \left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & 1 \\ \hline \end{array} \right) \cup Y \left(\begin{array}{|c|c|} \hline 2 & 1 \\ \hline 1 & 2 \\ \hline \end{array} \right) = T_2$$

Random Latin Squares Complexes

Multiple Latin Squares

For $\underline{L}^d = (L_1, \dots, L_d) \in \mathcal{L}_n^d$ let $Y(\underline{L}^d) = \cup_{i=1}^d Y(L_i)$.

The Probability Space $\mathcal{Y}(n, d)$

$\mathcal{L}_n^d = d$ -tuples of Latin squares of order n with uniform measure.

$\mathcal{Y}(n, d) = \{Y(\underline{L}^d) : \underline{L}^d \in \mathcal{L}_n^d\}$ with induced measure from \mathcal{L}_n^d .

Theorem [Lubotzky-M]:

There exist $\epsilon > 0, d < \infty$ such that

$$\lim_{n \rightarrow \infty} \Pr[Y \in \mathcal{Y}(n, d) : h_1(Y) > \epsilon] = 1.$$

Remark: $\epsilon = 10^{-11}$ and $d = 10^{11}$ will do.

Idea of Proof

Fix $0 < c < 1$ and let $\phi \in C^1(T_n; \mathbb{F}_2)$.

$$\phi \text{ is } \begin{cases} c - \text{small} & \text{if } \|\phi\| \leq cn^2 \\ c - \text{large} & \text{if } \|\phi\| \geq cn^2 \end{cases}$$

c -Small Cochains

Lower bound on expansion in terms of the spectral gap of the vertex links.

c -Large Cochains

Expansion is obtained by means of a new large deviations bound for the probability space \mathcal{L}_n of Latin squares.

2-Expansion and Spectral Gap

Notation

For a complex $T_n^{(1)} \subset Y \subset T_n$ let:

$Y_v = \text{lk}(Y, v) =$ the link of $v \in V$.

$\mu_v =$ spectral gap of the $n \times n$ bipartite graph Y_v .

$\tilde{\mu} = \min_{v \in V} \mu_v$.

$d = D_1(Y) =$ maximum edge degree in Y .

Theorem [LM]:

If $\|[\phi]\| \leq cn^2$ then

$$\|d_1\phi\| \geq \left(\frac{(1 - c^{1/3})\tilde{\mu}}{2} - \frac{d}{3} \right) \|[\phi]\|.$$

Spectral Gap of Random Graphs

Random Bipartite Graphs

$\tilde{\pi} = (\pi_1, \dots, \pi_d) \in \mathbb{S}_n^d$ defines a graph $G = G(\tilde{\pi})$ by

$$E(G) = \{ (i, \pi_j(i)) : 1 \leq i \leq n, 1 \leq j \leq d \} \subset [n]^2.$$

$\mathcal{G}(n, d)$ = uniform probability space $\{G(\tilde{\pi}) : \tilde{\pi} \in \mathbb{S}_n^d\}$.

Theorem [Friedman]:

For a fixed $d \geq 100$:

$$\Pr[G \in \mathcal{G}(n, d) : \mu_2(G) > d - 3\sqrt{d}] = 1 - O(n^{-2}).$$

Expansion of c -Small Cochains

Links as Random Graphs

Let $Y = Y(\underline{L}^d)$ be a random complex in $\mathcal{Y}(n, d)$.

Then $Y_\nu = \text{lk}(Y, \nu)$ is a random graph in $\mathcal{G}(n, d)$.

Therefore

$$\Pr[\tilde{\mu} \geq d - 3\sqrt{d}] = 1 - O(n^{-1}).$$

Corollary:

Let $d \geq 100$ and $c < 10^{-3}$. If $\|[\phi]\| \leq cn^2$ then

$$\begin{aligned} \frac{\|d_1\phi\|}{\|[\phi]\|} &\geq \frac{(1 - c^{1/3})\tilde{\mu}}{2} - \frac{d}{3} \\ &\geq \frac{(1 - c^{1/3})(d - 3\sqrt{d})}{2} - \frac{d}{3} > 1. \end{aligned}$$

Large Deviations for Latin Squares

The Random Variable $f_{\mathcal{E}}$

\mathcal{E} - a family of 2-simplices of T_n , $|\mathcal{E}| \geq cn^3$.

For a Latin square $L \in \mathcal{L}_n$ let

$$f_{\mathcal{E}}(L) = |Y(L) \cap \mathcal{E}|.$$

Then

$$E[f_{\mathcal{E}}] = \frac{|\mathcal{E}|}{n} \geq cn^2.$$

Theorem [LM]:

For all $n \geq n_0(c)$

$$\Pr[L \in \mathcal{L}_n : f_{\mathcal{E}}(L) < 10^{-3}c^2n^2] < e^{-10^{-3}c^2n^2}.$$

Topological Overlap Property for $\mathcal{Y}(n, d)$

Corollary:

There exist $\delta > 0$ and d such that $Y \in \mathcal{Y}(n, d)$ a.a.s. satisfies the following:

For any continuous map $f : Y \rightarrow \mathbb{R}^2$ there exists $p \in \mathbb{R}^2$ such that

$$\gamma_Y(p) \geq \delta n^2.$$

Lecture III: Garland's Method

Eigenvalues and Cohomology

- ▶ Higher Laplacians
- ▶ Simplicial Hodge theorem
- ▶ Garland's Theorem

Applications of Garland's method

- ▶ Cohomology of discrete groups
- ▶ Topology of random flag complexes
- ▶ Hypergraph matching and homology

Random Surfaces

- ▶ The Harer-Zagier model
- ▶ The genus of a random surface
- ▶ Related models

The Graph Laplacian

$G = (V, E)$ a graph

The **Laplacian** of G is the $V \times V$ matrix L_G :

$$L_G(u, v) = \begin{cases} \deg(u) & u = v \\ -1 & uv \in E \\ 0 & \text{otherwise} \end{cases}$$

Spectrum of L_G : $0 = \lambda_1(G) \leq \lambda_2(G) \leq \dots \leq \lambda_n(G)$

The Spectral Gap

$\lambda_2(G)$ controls the expansion of G and the convergence rate of a random walk on G . In particular, $\lambda_2(G) > 0 \Leftrightarrow G$ connected.

Higher Laplacians

A positive weight function $c(\sigma)$ on the simplices of X induces an **Inner product** on $C^k(X) = C^k(X; \mathbb{R})$:

$$(\phi, \psi) = \sum_{\sigma \in X(k)} c(\sigma) \phi(\sigma) \psi(\sigma) .$$

Adjoint $d_k^* : C^{k+1}(X) \rightarrow C^k(X)$

$$(d_k \phi, \psi) = (\phi, d_k^* \psi) .$$

$$C^{k-1}(X) \begin{array}{c} \xrightarrow{d_{k-1}} \\ \xleftarrow{d_{k-1}^*} \end{array} C^k(X) \begin{array}{c} \xrightarrow{d_k} \\ \xleftarrow{d_k^*} \end{array} C^{k+1}(X)$$

The reduced **k -Laplacian** of X is the positive semidefinite operator

$$\Delta_k = d_{k-1} d_{k-1}^* + d_k^* d_k : C^k(X) \rightarrow C^k(X) .$$

Harmonic Cochains

The space of **Harmonic** k -cochains

$$\ker \Delta_k = \{\phi \in C^k(X) : d_k \phi = 0, d_{k-1}^* \phi = 0\}.$$

Simplicial Hodge Theorem:

$$C^k(X) = \text{Im } d_{k-1} \oplus \ker \Delta_k \oplus \text{Im } d_k^* .$$

$$\ker \Delta_k \cong \tilde{H}^k(X; \mathbb{R}).$$

$\mu_k(X)$ = minimal eigenvalue of Δ_k .

A Vanishing Criterion:

$$\mu_k(X) > 0 \Leftrightarrow \tilde{H}^k(X; \mathbb{R}) = 0.$$

Eigenvalues and Cohomology

Let X be a pure d -dimensional complex with weight function:

$$c(\sigma) = (d - \dim \sigma)! |\{\tau \in X(d) : \tau \supset \sigma\}|.$$

For $\tau \in X$ consider the link $X_\tau = \text{lk}(X, \tau)$ with a weight function given by $c_\tau(\alpha) = c(\tau\alpha)$.

Theorem [Garland '72]:

Let $0 \leq \ell < k < d$. Then:

$$\min_{\tau \in X(\ell)} \mu_{k-\ell-1}(X_\tau) > \frac{\ell+1}{k+1} \Rightarrow H^k(X; \mathbb{R}) = 0.$$

In particular:

$$\min_{\tau \in X(d-2)} \mu_0(X_\tau) > \frac{d-1}{d} \Rightarrow H^{d-1}(X; \mathbb{R}) = 0.$$

Sketch of Proof I

For $\phi \in C^k(X)$ define $\phi_\tau \in C^{k-\ell-1}(X_\tau)$ by $\phi_\tau(\alpha) = \phi(\tau\alpha)$.

Garland's Identity

$$\binom{k}{\ell+1}(\Delta_k\phi, \phi) = \sum_{\tau \in X(\ell)} (\Delta_{k-\ell-1}\phi_\tau, \phi_\tau) - \binom{k}{\ell} \|\phi\|^2.$$

Proof of Garland's Theorem:

Suppose that

$$\min_{\tau \in X(\ell)} \mu_{k-\ell-1}(X_\tau) > \frac{\ell+1}{k+1}$$

and let $0 \neq \phi \in C^k(X)$ such that $\Delta_k\phi = \mu_k(X)\phi$.

Sketch of Proof II

By Garland's identity:

$$\begin{aligned}\mu_k(X) \binom{k}{\ell+1} \|\phi\|^2 &= \binom{k}{\ell+1} (\Delta_k \phi, \phi) \\ &= \sum_{\tau \in X(\ell)} (\Delta_{k-\ell-1} \phi_\tau, \phi_\tau) - \binom{k}{\ell} \|\phi\|^2 \\ &\geq \min_{\tau \in X(\ell)} \mu_{k-\ell-1}(X_\tau) \sum_{\tau \in X(\ell)} \|\phi_\tau\|^2 - \binom{k}{\ell} \|\phi\|^2 \\ &\geq \left(\min_{\tau \in X(\ell)} \mu_{k-\ell-1}(X_\tau) \binom{k+1}{\ell+1} - \binom{k}{\ell} \right) \|\phi\|^2 \\ &> \left(\frac{\ell+1}{k+1} \binom{k+1}{\ell+1} - \binom{k}{\ell} \right) \|\phi\|^2 = 0.\end{aligned}$$

Complexes with Expanding Links

The Projective Plane Graph

$G_q = (V_q, E_q)$: points vs. lines graph of $PG(2, q)$.

$$|V_q| = 2(q^2 + q + 1) \quad , \quad |E_q| = (q + 1)(q^2 + q + 1).$$

Spectral Gap: $\mu_0(G_q) = 1 - \frac{\sqrt{q}}{q+1}$.

If $q \geq d^2$ then $\mu_0(G_q) > \frac{d-1}{d}$. This implies the following

Theorem [Garland]:

Let $q \geq d^2$ and let X be a pure d -dimensional complex such that $\text{lk}(X, \tau) \cong G_q$ for all $\tau \in X(d-2)$.

Then $H_{d-1}(X; \mathbb{R}) = 0$.

Cohomology of Discrete Subgroups

\mathbb{K} a local field with residue field \mathbb{F}_q .

Γ a torsion-free discrete cocompact subgroup of $SL_{d+1}(\mathbb{K})$.

Theorem [Garland]:

If $q \geq d^2$ then $H_{d-1}(\Gamma, \mathbb{R}) = 0$.

Sketch of Proof:

$\mathcal{B} = \tilde{A}_d(\mathbb{K})$ - the affine building associated to $SL_{d+1}(\mathbb{K})$.

\mathcal{B} is a contractible complex with a free Γ action.


The quotient space $B\Gamma = \mathcal{B}/\Gamma$ is a pure d -dimensional complex such that $\text{lk}(B\Gamma, \tau) \cong G_q$ for all $\tau \in B\Gamma(d-2)$.

Therefore


$$H^{d-1}(\Gamma; \mathbb{R}) = H^{d-1}(B\Gamma; \mathbb{R}) = 0.$$

Flag Complexes

The **flag complex** $X(G)$ of a graph $G = (V, E)$:
Vertex set: V , Simplices: all cliques σ of G .



G



$X(G)$


Remarks:

The first subdivision of a complex is a flag complex.

Face Numbers of Flag Complexes

Octahedral n -Sphere

$$(S^0)^{*(k+1)} = \{a_1, b_1\} * \cdots * \{a_{k+1}, b_{k+1}\}$$



Proposition [M '03]:

If $\tilde{H}_k(X(G)) \neq 0$ then for all j :

$$f_j(X(G)) \geq f_j((S^0)^{*(k+1)}) = \binom{k+1}{j+1} 2^{j+1}.$$

Homology of Flag Complexes of Random Graphs

Let $\epsilon > 0$ be fixed and let $G \in G(n, p)$.

Theorem [Kahle '12]:

$$p \leq n^{-\frac{1}{k}-\epsilon} \Rightarrow H_k(X(G); \mathbb{Z}) = 0 \text{ a.a.s.}$$

$$p \geq \left(\frac{(\frac{k}{2} + 1 + \epsilon) \log n}{n} \right)^{\frac{1}{k+1}} \Rightarrow H_k(X(G); \mathbb{R}) = 0 \text{ a.a.s.}$$

Theorem [DeMarco-Hamm-Kahn '12]:

$$p \geq \left(\frac{(\frac{3}{2} + \epsilon) \log n}{n} \right)^{\frac{1}{2}} \Rightarrow H_1(X(G); \mathbb{F}_2) = 0 \text{ a.a.s.}$$

Vanishing of $H_k(X(G); \mathbb{R})$

Let $C_k = \frac{k}{2} + 1 + \epsilon$ and let $p = \left(\frac{C_k \log n}{n}\right)^{\frac{1}{k+1}}$.

Claim 1: The $(k+1)$ -skeleton $Y = X(G)^{(k+1)}$ is a.a.s. pure:

$$\begin{aligned} & E[\#\sigma \in Y(k) \text{ such that } \sigma \not\subset (k+1)\text{-face of } Y] \\ &= \binom{n}{k+1} p^{\binom{k+1}{2}} (1 - p^{k+1})^{n-k-1} \\ &\leq n^{k+1} \left(\frac{C_k \log n}{n}\right)^{\frac{k}{2}} \left(1 - \frac{C_k \log n}{n}\right)^{n-k-1} = O(n^{-\frac{\epsilon}{2}}). \end{aligned}$$

Claim 2: $\mu_0(\text{lk}(Y, \tau)) = 1 - o(1)$ a.a.s. for all $\tau \in Y(k-1)$.

Claims 1 & 2 + Garland's Thm. $\Rightarrow H_k(X(G); \mathbb{R}) = 0$ a.a.s.

Fundamental Group of $X(G)$


Theorem [Babson]:

If $n^{-\frac{1}{2}+\epsilon} < p < n^{-\frac{1}{3}-\epsilon}$ then a.a.s. $\pi_1(X(G))$ is nontrivial and hyperbolic.

Theorem [Costa-Farber-Horak]:

- ▶ If $n^{-\frac{11}{30}+\epsilon} < p < n^{-\frac{1}{3}-\epsilon}$ then a.a.s. $\pi_1(Y)$ has 2-torsion.
- ▶ If $n^{-\frac{1}{2}+\epsilon} < p < n^{-\frac{11}{30}-\epsilon}$ then a.a.s. $\pi_1(Y)$ is torsion free.

Flag Projective Plane



If $p = n^{-\frac{f_0}{f_1}+\epsilon} = n^{-\frac{11}{30}+\epsilon}$ then
a.a.s. $X(G) \supset Z \cong \mathbb{P}^2$ such that
 $\mathbb{Z}_2 = \pi_1(Z) \hookrightarrow \pi_1(Y)$.

Spectral Gap and Homological Connectivity

Theorem [Aharoni-Berger-M]:

$G = (V, E)$ - a graph, $|V| = n$.

$$\lambda_2(G) > \frac{(k-1)n}{k} \Rightarrow \tilde{H}_i(X(G); \mathbb{R}) = 0 \quad \forall 0 \leq i \leq k-1.$$

The Bound is Sharp

$n = k\ell$.

$T_k(n)$ - the complete k -partite graph with equal sides.

$$\lambda_2(T_k(n)) = \frac{(k-1)n}{k}, \quad \dim \tilde{H}_{k-1}(X(T_k(n)); \mathbb{R}) = (\ell-1)^k \neq 0.$$

The theorem has applications to hypergraph matching ...

Bipartite Matching

A_1, \dots, A_m finite sets.

A **System of Distinct Representatives (SDR)**:

a choice of **distinct** $x_1 \in A_1, \dots, x_m \in A_m$.

A_1	A_2	A_3
1	1	
		2
3	3	3

\exists SDR

A_1	A_2	A_3
1	1	
2		2

\nexists SDR

Hall's Theorem (1935)

(A_1, \dots, A_m) has an SDR iff

$|\cup_{i \in I} A_i| \geq |I|$ for all $I \subset [m] = \{1, \dots, m\}$.

Hypergraph Matching

A **Hypergraph** is a family of sets $\mathcal{F} \subset 2^V$

$(\mathcal{F}_1, \dots, \mathcal{F}_m)$ a sequence of m hypergraphs

A **System of Disjoint Representatives (SDR)** for $(\mathcal{F}_1, \dots, \mathcal{F}_m)$


is a choice of **pairwise disjoint** $F_1 \in \mathcal{F}_1, \dots, F_m \in \mathcal{F}_m$

When do $(\mathcal{F}_1, \dots, \mathcal{F}_m)$ have an SDR?


The problem is **NP-Complete** even if all \mathcal{F}_i 's consist of 2-element sets. Therefore, we cannot expect a "good" characterization as in Hall's Theorem.

There are however some interesting sufficient conditions ...




Do $(\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3, \mathcal{F}_4)$ have an SDR?



\exists SDR



\nexists SDR




The Aharoni-Haxell Theorem


A **Matching** is a hypergraph \mathcal{M} of **pairwise disjoint** sets.

The **Matching Number** $\nu(\mathcal{F})$ of a hypergraph \mathcal{F} is the maximal size $|\mathcal{M}|$ of a matching $\mathcal{M} \subset \mathcal{F}$.

$$\nu(\mathcal{F}) = 3$$



$$\nu(\mathcal{F}) = 1$$



The Aharoni-Haxell Theorem

$\mathcal{F}_1, \dots, \mathcal{F}_m \subset \binom{V}{r}$ such that for all $I \subset [m]$

$$\nu\left(\bigcup_{i \in I} \mathcal{F}_i\right) > r(|I| - 1) .$$

Then $(\mathcal{F}_1, \dots, \mathcal{F}_m)$ has an SDR.

A Fractional Extension

A **Fractional Matching** of a hypergraph \mathcal{F} on V is a function $f : \mathcal{F} \rightarrow \mathbb{R}_+$ such that $\sum_{F \ni v} f(F) \leq 1$ for all $v \in V$.

The **Fractional Matching Number** $\nu^*(\mathcal{F})$ is $\max_f \sum_{F \in \mathcal{F}} f(F)$ over all fractional matchings f .

Example: The Finite Projective Plane \mathcal{P}_n

$$\nu(\mathcal{P}_n) = 1 \quad , \quad \nu^*(\mathcal{P}_n) = \frac{n^2+n+1}{n+1}$$

Theorem [ABM]:

$\mathcal{F}_1, \dots, \mathcal{F}_m \subset \binom{V}{r}$ such that for all $I \subset [m]$

$$\nu^*\left(\bigcup_{i \in I} \mathcal{F}_i\right) > r(|I| - 1) \quad .$$

Then $(\mathcal{F}_1, \dots, \mathcal{F}_m)$ has an SDR.

Random Orientable Surfaces

Harer-Zagier Model

P - a regular $2n$ -gon with vertices $1, \dots, 2n$ and edges

$e_1 = (2n \rightarrow 1)$, $e_2 = (1 \rightarrow 2)$, \dots , $e_{2n} = (2n - 1 \rightarrow 2n)$.

C_{2n} - the conjugacy class of fixed point free involutions in S_{2n} .

$|C_{2n}| = (2n - 1)!! = (2n - 1) \cdot (2n - 3) \cdots 3 \cdot 1$.

For $\pi \in C_{2n}$ glue all pairs of edges $e_i, e_{\pi(i)}$ with reverse orientations to obtain an orientable surface $M(\pi)$.

Genus of $M(\pi)$

$N(\eta)$ = the number of cycles in $\eta \in S_{2n}$.


$\sigma = (123 \cdots 2n) \in S_{2n}$.

Fact: $f_0(M(\pi)) = N(\pi\sigma)$.

Therefore:


$$2 - 2 \cdot \text{genus}(M(\pi)) = f_0 - f_1 + f_2 = N(\pi\sigma) - n + 1.$$

Example: A Sphere




$$g = \frac{n+1 - N(\pi\sigma)}{2} = \frac{2+1-3}{2} = 0.$$

Example: A Torus




$$\xrightarrow[\pi\sigma = (13)(24)(1234) = (1432)]{\pi = (13)(24)}$$



$$g = \frac{n+1 - N(\pi\sigma)}{2} = \frac{2+1-1}{2} = 1.$$

Example: A Genus 2 Surface



$$g = \frac{n+1 - N(\pi\sigma)}{2} = \frac{4+1-1}{2} = 2.$$

Distribution of the Genus

Harer-Zagier Formula

Let $p_n(g) = \Pr[\pi \in C_{2n} : \text{genus}(M(\pi)) = g]$.

Then:

$$1 + 2 \sum_{n,g} p_n(g) k^{n+1-2g} x^{n+1} = \left(\frac{1+x}{1-x} \right)^k.$$

Expectation and Variance

$$E[\text{genus}] = \frac{1}{2}(n+1 - \log(2n) - \gamma) + O\left(\frac{1}{n}\right).$$

$$\text{Var}[\text{genus}] = \frac{1}{4}(\log(2n) + \gamma - \frac{\pi^2}{6}) + O\left(\frac{\log n}{n}\right).$$